

Conference Abstract

Science and Cycling Conference, Barcelona 2026

Acute Hemodynamic Responses to Three Work-Matched HIIT Training Protocols in Trained Cyclists

Görkem A. Balci ^{1*}, Hakan Arslan ¹, Burak A. Ünsal ¹, Ramazan Aydınoğlu ¹ and Bent R. Rønnestad ²

Received: 2 March 2026

Accepted: 24 March 2026

Published: 29 October 2026

¹ Coaching Education Department, Faculty of Sports Sciences, Ege University, İzmir, Türkiye² Section for Health and Exercise Physiology, University of Inland Norway, Lillehammer, Norway

Correspondence

Görkem A. Balci

Coaching Education Department, Faculty of Sports Sciences, Ege University, İzmir, Türkiye

gorkem.aybars.balci@ege.edu.tr

Abstract

Current literature indicates that studies comprehensively examining acute central and peripheral hemodynamic responses to different high-intensity interval training (HIIT) modalities in trained cyclists remain limited. Therefore, this study aims to compare the acute central cardiac output (Q), stroke volume (SV) and peripheral arteriovenous O₂ difference (a-vO_{2diff}) responses elicited by three distinct HIIT models and to determine which modality provides a greater acute stimulus to the respective physiological components. Twelve trained male cyclists (Level 3-4) participated in this study (age: 29.7 ± 5.3 years; maximal oxygen uptake (VO_{2max}): 57.8 ± 4.1 mL·min⁻¹·kg⁻¹; body mass: 69.8 ± 6.0 kg). Acute hemodynamic responses were evaluated with three work- and effort-matched HIIT models administered in a randomised order: Long-HIIT (4×3 min work- 3 min recoveries), Short-HIIT (1 series of 15×43.5 sec work - 43.5 sec recovery), and Decreasing Power Output HIIT (DPO-HIIT; 6×114.5 sec work - 114.5 sec recovery). Q, SV, and a-vO_{2diff} were measured using a non-invasive inert gas rebreathing method (N₂ORB). Two-way repeated-measures analysis of variance revealed no significant main effect of the protocol for Q (F(2.000, 22.000)=1.966, p=0.178, η²=0.152), SV (F(2.000, 22.000)=0.790, p=0.446, η²=0.067), heart rate (HR) (F(2.000, 22.000)=0.620, p=0.547, η²=0.053), or a-vO_{2diff} (F(2.000, 22.000)=1.586, p=0.227, η²=0.126). The Protocol × Time interaction was significant only for HR (F(2.467, 27.132)=4.561, p=0.014, η²=0.293). Consequently, while overall systemic stress remains matched, coaches and athletes can utilise these different HIIT structures interchangeably, which may allow for flexibility in endurance training prescription.

Keywords

arterio-venous O₂ difference; high-intensity interval training; cardiac output; performance; stroke volume

1 Introduction

Cardiac output (Q), stroke volume (SV), and the arteriovenous O_2 difference ($a-vO_{2diff}$) together constitute the primary determinants of maximal oxygen uptake ($\dot{V}O_{2max}$), which is a critical physiological factor determining endurance performance (Rønnestad et al., 2020). SV and heart rate (HR), which together determine Q , represent the central component of $\dot{V}O_{2max}$, whereas the $a-vO_{2diff}$ constitutes the peripheral component (Faricier et al., 2025; Joyner & Dominelli, 2021). Individual differences in physiological and haemodynamic responses should therefore be considered when targeting cardiovascular and peripheral adaptations through HIIT (Buchheit & Laursen, 2013). Accordingly, for an athlete with a larger deficit in the central component, high-intensity interval training (HIIT) protocols that elicit and sustain higher peak Q and/or peak SV for longer durations may be considerably more effective. Current literature indicates that studies comprehensively examining acute central and peripheral hemodynamic responses to different HIIT modalities in trained cyclists remain limited. Therefore, this study aims to compare the acute central (Q or SV) and peripheral ($a-vO_{2diff}$) responses elicited by three distinct HIIT models and to determine which modality provides a greater acute stimulus to the respective physiological components.

2 Material and Methods

2.1 Participants

Twelve trained male cyclists (Level 3-4), (De Pauw et al., 2013) participated in this study (age: 29.7 ± 5.3 years; $\dot{V}O_{2max}$: 57.8 ± 4.1 mL $\cdot$ min $^{-1}\cdot$ kg $^{-1}$; body mass: 69.8 ± 6.0 kg). The study was conducted in accordance with the Declaration of Helsinki and received institutional ethics committee approval (Decision No: 24-11.1T/41; Date: 28 November 2024).

2.2 Design

A randomised repeated-measures design was used in this study. Acute hemodynamic responses were evaluated with three work- and effort-matched ($RPE \geq 19$) HIIT models administered in a randomized order: Long HIT (4 $\times$ 3 min work - 3 min recovery; $WR \sim 88\%$ of peak WR (WR_{peak}); Recovery (REC) $\sim 25\%$ of WR_{peak}), Short-HIIT (1 series of 15 $\times$ 43.5 sec work - 43.5 sec recovery; $WR \sim 110\%$ of WR_{peak} ; REC $\sim 10\%$ of WR_{peak}) and Decreasing Power Output HIIT (DPO-HIIT, 6 $\times$ 114.5 sec work - 114.5 sec recovery; WR starting at I_{HIGH} and decreased by 4 equal steps to the $CP + 10\%$; Rec $\sim 25\%$ of WR_{peak}).

2.3 Methodology

Familiarisation sessions were conducted to adapt the participants to the electromagnetically braked cycle ergometer (Lode Excalibur Sport, Groningen, Netherlands) and cardiac measurement system (Innocor, COSMED, Italy).

$\dot{V}O_{2max}$ was determined via a ramp incremental test, with the WR increasing by 1 W every 2 s (30 W $\cdot$ min $^{-1}$).

Critical power and the highest constant-intensity that permits the attainment of $\dot{V}O_{2max}$ (I_{HIGH}) were determined using multiple constant-work-rate testing sessions (Caputo & Denadai, 2008; Jones et al., 2010).

Q , SV , and $a-vO_{2diff}$ were measured using a valid and reliable non-invasive inert gas rebreathing method (N_2O_{RB}), (Fontana et al., 2009).

Throughout the sessions, variables were recorded at four time points (Δ Time: 25%, 50%, 75%, and 100% of each bout).

2.4 Statistical Analysis

Data were analysed using a two-way (Protocol $\times$ Δ Time) repeated-measures ANOVA with Greenhouse-Geisser corrections

where appropriate. Partial eta squared (η^2) was reported as the effect size for the overall effect of protocol in the repeated-measures ANOVA and was interpreted as small ($\eta^2 \geq 0.01$), medium ($\eta^2 \geq 0.06$), and large ($\eta^2 \geq 0.14$) (Richardson, 2011) (Richardson, 2011).

Bonferroni-adjusted post-hoc tests were applied for pairwise comparisons. Cohen's d was reported for pairwise post-hoc comparisons to quantify the magnitude of standardised differences between individual HIIT protocols and was interpreted as small ($d \geq 0.20$), medium ($d \geq 0.50$), and large ($d \geq 0.80$) (Cohen, 1988). Statistical analyses were performed using JASP (Version 0.98.1; JASP Team, 2026).

3 Results

Two-way repeated-measures ANOVA revealed no significant main effect of protocol for Q ($F(2.000, 22.000) = 1.966$, $p = 0.178$, $\eta^2 = 0.152$), SV ($F(2.000, 22.000) = 0.790$, $p = 0.446$, $\eta^2 = 0.067$), HR ($F(2.000, 22.000) = 0.620$, $p = 0.547$, $\eta^2 = 0.053$), or $a-vO_{2\text{diff}}$ ($F(2.000, 22.000) = 1.586$, $p = 0.227$, $\eta^2 = 0.126$). The Protocol $\times$ Δ Time interaction was significant only for HR ($F(2.467, 27.132) = 4.561$, $p = 0.014$, $\eta^2 = 0.293$). The Protocol $\times$ Δ Time interactions for Q ($F(2.476, 27.232) = 2.448$, $p = 0.095$, $\eta^2 = 0.182$), SV ($F(2.330, 25.629) = 0.620$, $p = 0.569$, $\eta^2 = 0.053$), and $a-vO_{2\text{diff}}$ ($F(3.043, 33.477) = 2.314$, $p = 0.093$, $\eta^2 = 0.174$) were not significant. The Protocol $\times$ Δ Time effect sizes for Q and $a-vO_{2\text{diff}}$ were large. The temporal responses of Q, SV, $a-vO_{2\text{diff}}$, and HR across the three HIIT protocols are presented in Figure 1.




Figure 1. Acute hemodynamic responses during the three high-intensity interval training protocols: (a) cardiac output (Q); (b) stroke volume (SV); (c) arteriovenous O₂ difference ($a-vO_{2\text{diff}}$); and (d) heart rate (HR) at Δ25, Δ50, Δ75, and Δ100 of each bout. Data are presented as mean $\pm$ SD.

Q increased over time across all protocols (Figure 1a). Bonferroni-adjusted pairwise comparisons, averaged across protocols, showed an increase from $\Delta 25$ to $\Delta 100$ ($p = 0.001$).

SV showed no significant change from $\Delta 25$ to $\Delta 100$ when averaged across protocols (Figure 1b). The Bonferroni-adjusted pairwise comparison between $\Delta 25$ and $\Delta 100$ was not significant ($p = 0.187$).

The $a-vO_{2diff}$ showed no significant change from $\Delta 25$ to $\Delta 100$ when averaged across protocols (Figure 1c). The Bonferroni-adjusted pairwise comparison between $\Delta 25$ and $\Delta 100$ was not different ($p = 0.415$).

HR increased significantly from $\Delta 25$ to $\Delta 100$ across protocols (Figure 1d). No significant main effect of protocol was observed, indicating that overall HR responses were similar between HIIT protocols. However, the significant Protocol $\times$ Δ Time interaction suggests that the temporal pattern of HR increase was not identical across protocols.

4 Discussion

The absence of a main protocol effect for Q, SV, and $a-vO_{2diff}$, further supported by trivial effect sizes in pairwise comparisons (Cohen's $d < 0.2$), aligns with the observation that, despite differences in interval structure, the three work-matched models elicited broadly comparable average central and peripheral responses.

In our data, Q increased over time, while SV remained stable. Therefore, the increase in Q seems to be mainly related to the increase in HR. For $a-vO_{2diff}$, we did not observe a clear protocol-specific difference. This pattern suggests that central hemodynamic adjustments contributed more to the temporal rise in oxygen transport than peripheral extraction.

Previous studies provide useful but different perspectives. Stanley and Buchheit (2014) examined acute cardiovascular responses during HIIT in well-trained cyclists and reported that SV could be maintained during interval exercise, regardless of recovery intensity (Stanley & Buchheit, 2014). This is consistent with our finding that SV remained relatively stable across the three HIIT protocols.

Coe and Astorino (2024) compared different HIIT formats and showed that protocol structure can influence acute hemodynamic responses such as Q and SV (Coe & Astorino, 2024). However, their protocols differed in total duration and exercise volume. Therefore, it is difficult to separate the effect of interval structure from the effect of total work or duration.

In contrast, our study matched total work, total time, and effort across protocols. So, our design provides a more controlled comparison of how interval structure itself may influence acute central and peripheral hemodynamic responses.

Although the Protocol $\times$ Δ Time interactions for Q and $a-vO_{2diff}$ exhibited large effect sizes ($\eta^2 = 0.182$ and 0.174 , respectively), subsequent pairwise comparisons revealed only trivial effect sizes (Cohen's $d < 0.2$) between the HIIT protocols. Therefore, the practical superiority of one specific HIIT structure over another regarding acute central and peripheral hemodynamic responses cannot be conclusively established from the current data.

5 Practical Applications

Consequently, while overall systemic stress remains matched across the different HIIT intervals, manipulating the specific structure of work-matched HIIT (Long, Short, or DPO) does not appear to selectively target central or peripheral oxygen extraction kinetics. Based on

our data, we cannot make protocol-specific recommendations for cyclists with this training status. Overall, these findings should be interpreted as practical tendencies rather than clear evidence of protocol superiority.

6 Conclusions

When overall work and effort are strictly matched, different HIIT structures (Long, Short, and DPO) elicit comparable acute central ($\dot{V}O_2$, SV) and peripheral ($a-vO_{2\text{diff}}$) hemodynamic stress. Therefore, manipulating work-effort match interval structure does not selectively isolate specific central or peripheral physiological responses. This may provide coaches and athletes with the flexibility to utilise any of these formats interchangeably to achieve a similar overall cardiovascular stimulus.

Funding: This study was supported by The Scientific and Technological Research Council of Türkiye (TÜBİTAK) under the 1001 Scientific and Technological Research Projects Funding Program (Project No: 224S942).

Acknowledgements: We would like to thank all participants for their time, effort, and commitment throughout the study. We also sincerely appreciate their cooperation during the laboratory testing sessions.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

- Buchheit, M., & Laursen, P. B. (2013). High Intensity Interval Training, Solutions to the Programming Puzzle: Part I: Cardiopulmonary Emphasis. *Sports Medicine*, 43(5), 313–338. <https://doi.org/10.1007/s40279-013-0029-x>
- Caputo, F., & Denadai, B. (2008). The highest intensity and the shortest duration permitting attainment of maximal oxygen uptake during cycling: Effects of different methods and aerobic fitness level. *European Journal of Applied Physiology*, 103, 47–57. <https://doi.org/10.1007/s00421-008-0670-5>
- Coe, L. N., & Astorino, T. A. (2024). Sex differences in hemodynamic response to high-intensity interval exercise. *Scandinavian Journal of Medicine & Science in Sports*, 34(1), e14495. <https://doi.org/10.1111/sms.14495>
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Lawrence Erlbaum Associates. <https://doi.org/10.4324/9780203771587>
- De Pauw, K., Roelands, B., Cheung, S., de Geus, B., Rietjens, G., & Meeusen, R. (2013). Guidelines to Classify Subject Groups in Sport-Science Research. *International Journal of Sports Physiology and Performance*, 8, 111–122. <https://doi.org/10.1123/ijsp.8.2.111>
- Faricier, R., Paterson, D., & Murias, J. (2025). Physiological Determinants of $\dot{V}O_{2\text{max}}$ Increase with Endurance Training in a Group Including Older and Young Adults. *Medicine & Science in Sports & Exercise*, 57. <https://doi.org/10.1249/MSS.0000000000003707>
- Fontana, P., Boutellier, U., & Toigo, M. (2009). Reliability of Measurements with InnocorTM during Exercise. *International Journal of Sports Medicine*, 30, 747–753. <https://doi.org/10.1055/s-0029-1225340>
- Jones, A., Vanhatalo, A., Burnley, M., Morton, R., & Poole, D. (2010). *Critical Power: Implications for the Determination of $\dot{V}O_{2\text{max}}$ and Exercise Tolerance*.
- Joyner, M. J., & Dominelli, P. B. (2021). Central cardiovascular system limits to aerobic capacity. *Experimental Physiology*, 106(12), 2299–2303. <https://doi.org/10.1113/EP088187>
- Richardson, J. T. E. (2011). Eta squared and partial eta squared as measures of effect size in educational research. *Educational Research Review*, 6(2), 135–147. <https://doi.org/10.1016/j.edurev.2010.12.001>
- Rønnestad, B. R., Hansen, J., Nygaard, H., & Lundby, C. (2020). Superior performance improvements in elite cyclists following short-interval vs effort-matched long-interval training. *Scandinavian Journal of Medicine & Science in Sports*, 30(5), 849–857. <https://doi.org/10.1111/sms.13627>
- Stanley, J., & Buchheit, M. (2014). Moderate Recovery Unnecessary to Sustain High Stroke Volume during Interval Training. A Brief Report. *Journal of sports science & medicine*, 13(2), 393–396.